

University of Nebraska - Lincoln

DigitalCommons@University of Nebraska - Lincoln

Agronomy & Horticulture -- Faculty Publications

Agronomy and Horticulture Department

2020

A justification for continued management of turfgrass during economic contraction

James T. Brosnan

The Univeristy of Tennessee, jbrosnan@utk.edu

Ambika Chandra

Texas A&M AgriLife Research and Extension Center

Roch E. Gaussoin

University of Nebraska - Lincoln, rgaussoin1@unl.edu

Alec Kowalewski

Oregon State University

Bernd Leinauer

Extension Plant Sciences, N Skeen Hall

See next page for additional authors

Follow this and additional works at: <https://digitalcommons.unl.edu/agronomyfacpub>



Part of the [Agricultural Science Commons](#), [Agriculture Commons](#), [Agronomy and Crop Sciences Commons](#), [Botany Commons](#), [Horticulture Commons](#), [Other Plant Sciences Commons](#), and the [Plant Biology Commons](#)

Brosnan, James T.; Chandra, Ambika; Gaussoin, Roch E.; Kowalewski, Alec; Leinauer, Bernd; Rossi, Frank S.; Soldat, Douglas J.; Stier, John C.; and Unruh, J. Bryan, "A justification for continued management of turfgrass during economic contraction" (2020). *Agronomy & Horticulture -- Faculty Publications*. 1390. <https://digitalcommons.unl.edu/agronomyfacpub/1390>






This Article is brought to you for free and open access by the Agronomy and Horticulture Department at DigitalCommons@University of Nebraska - Lincoln. It has been accepted for inclusion in Agronomy & Horticulture -- Faculty Publications by an authorized administrator of DigitalCommons@University of Nebraska - Lincoln.

Authors

James T. Brosnan, Ambika Chandra, Roch E. Gaussoin, Alec Kowalewski, Bernd Leinauer, Frank S. Rossi, Douglas J. Soldat, John C. Stier, and J. Bryan Unruh

COMMENTARY

A justification for continued management of turfgrass during economic contraction

James T. Brosnan¹  | Ambika Chandra² | Roch E. Gaussoin³ |
Alec Kowalewski⁴  | Bernd Leinauer⁵  | Frank S. Rossi⁶ | Douglas J. Soldat⁷ |
John C. Stier¹  | J. Bryan Unruh⁸ 

¹ Dep. of Plant Sciences, The Univ. of Tennessee, 112 Plant Biotechnology Building, 2505 EJ Chapman Dr., Knoxville, TN 37996, USA

² Texas A&M AgriLife Research and Extension Center, 17360 Coit Rd., Dallas, TX 75252, USA

³ Dep. of Agronomy and Horticulture, Univ. of Nebraska-Lincoln, 219 Plant Science Hall, Lincoln, NE 68583, USA

⁴ Dep. of Horticulture, Oregon State Univ., 4147 ALS Building, Corvallis, OR 97331, USA

⁵ Extension Plant Sciences, N Skeen Hall, Las Cruces, NM 88003, USA

⁶ Cornell Univ. School of Integrative Plant Science, G47 Plant Science Building, Ithaca, NY 14853, USA

⁷ Dep. of Soil Science Univ. of Wisconsin-Madison, 1525 Observatory Dr., Madison, WI 53706, USA

⁸ West Florida Research and Education Center, 4253 Experiment Dr., Hwy. 182, Jay, FL 32565, USA

Correspondence

James T. Brosnan, Dep. of Plant Sciences, The Univ. of Tennessee, 112 Plant Biotechnology Building, 2505 EJ Chapman Dr., Knoxville, TN 37996, USA.
Email: jbrosnan@utk.edu

Abstract

A novel coronavirus, termed COVID-19, spread worldwide and become a global pandemic in 2020. Forecasts show that COVID-19 will cause substantial economic contraction affecting almost every industry. Managed turfgrass, particularly in urban settings, has many positive societal and environmental benefits. In a contracted economy, will resources be available to manage turfgrass to achieve these benefits? In this paper, we outline the benefits of managed turfgrass on golf courses, playing fields, recreational parks, and urban landscapes to assist decision makers with resource allocation in the COVID-19 era.

1 | INTRODUCTION

A novel coronavirus, termed COVID-19 by the World Health Organization, rapidly spread worldwide and become a global pandemic (Sohrabi et al., 2020). By

This is an open access article under the terms of the [Creative Commons Attribution-NonCommercial-NoDerivs](https://creativecommons.org/licenses/by-nc-nd/4.0/) License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

© 2020 The Authors. *Agricultural & Environmental Letters* published by Wiley Periodicals LLC on behalf of American Society of Agronomy, Crop Science Society of America, and Soil Science Society of America

4 Aug. 2020, there were nearly 18.1 million confirmed cases worldwide, including 4.6 million in the United States leading to more than 154,000 deaths (WHO 2020). Travel bans, enforced quarantines, school closures, and social distancing practices were implemented to limit spread of COVID-19 (Parmet and Sinha, 2020; Sohrabi et al., 2020). The National Bureau of Economics Research estimates that COVID-19 will lead to a 20% contraction of the U.S. economy by the end of the fiscal year, with effects continuing into the future as businesses reduce expenditures related to innovation, training, and general management (Baker, Bloom, Davis, & Terry, 2020).

Nearly every industry, including turfgrass management, has been affected by COVID-19. In addition to the cancellation or delay of professional and collegiate sports played on turfgrass surfaces (PGA Tour, Major League Baseball, Major League Soccer, and National Collegiate Athletic Association), COVID-19 led to the temporary closure of all golf courses in 19 states (NGF, 2020) as well as municipal parks and recreational fields across the United States. In some states, lawn care and outdoor activities were restricted or prohibited (State of Michigan, 2020; State of Oregon, 2020). Many facilities have furloughed staff, and it remains uncertain if financial resources will be available to fill these positions in the future. A federal order facilitated the landscape industry to maintain normal staffing during the pandemic (Krebs, 2020), but this may change with the predicted downturn of the economy.

COVID-19-induced economic uncertainty will have a major effect on all facets of agriculture for the foreseeable future (Lusk et al., 2020). It is well established that managed turfgrass, particularly in urban settings, offers many societal and environmental benefits. In a contracted economy, will resources be available to manage turfgrass to achieve these benefits? Our objective here is to outline the benefits of managed turfgrass on golf courses, playing fields, recreational parks, and urban landscapes to assist decision makers with resource allocation in the COVID-19 era.

2 | BENEFITS OF TURFGRASS

Turfgrasses ameliorate the impacts of urban sprawl in today's rapidly urbanizing global society. The many environmental benefits offered have been documented by several authors and summarized by Beard and Green (1994), Stier et al. (2013), and Christians, Patton, and Law (2017). Benefits include temperature moderation through evaporative cooling resulting in dissipation of radiant heat and mitigation of the heat island effect in cities (Amani-Beni, Zhang, Gao-di, & Jie, 2018; Jenerette, Harlan, Stefanov, & Martin, 2011; Wang et al., 2016b).

Core Ideas

- Managed turfgrass confers many positive societal and environmental benefits.
- These ecosystem services are predicated on sites receiving appropriate management resources.
- Resource allocation in the COVID-19 era should be structured to achieve these benefits.

Maximum daily canopy temperature of a green, growing turfgrass was found to be 21 °C cooler than a brown, dormant, sward and 39 °C cooler than synthetic turf (Beard & Johns, 1985). The transpirational cooling effect of green turfgrass can reduce energy input and costs required for indoor cooling (Beard & Beard, 1985). Turfgrasses also play an important role in stabilizing soils in urban centers, on underutilized and abandoned properties (Montgomery, et al, 2016); reducing surface water runoff, thereby recharging groundwater and reducing soil and water erosion (Bouwer, 2002; Dai, Puyang, & Han, 2016); absorbing atmospheric pollutants (greenhouse gases) produced through anthropogenic activities, especially in densely populated regions (Thwaites et al., 2006), sequestering carbon (Braun & Bremer, 2019; Hamido, Guertal, & Wood, 2016; Law and Patton, 2017; Qian and Follett, 2002), producing oxygen, and improving air quality (Monteiro, 2017); reducing noise pollution (Van Renterghem et al., 2015); and decontaminating soils through phytoremediation and associated plant-microbe interactions (Crouch, Carter, Ismaiel, & Roberts, 2017; Grant, Campbell, & Charnock, 2002; Krishnan et al., 2000; Lin et al., 2004; Shahandeh and Hossner, 2000; Thompson et al., 2008; Tordoff, 2000).

Turfgrasses also positively affect people by improving their physical and mental health and social well-being. Greenspace, which includes healthy lawns and landscapes, is inversely related to the incidence of crime, especially in urban neighborhoods (Kuo and Sullivan, 2001; Kuo et al., 1998). A well-managed lawn with healthy turfgrass, trees, and ornamental plants increases property value by improving perceptions of residential communities, businesses, and schools (Laverne and Winsin-Geideman, 2003; Stier et al., 2013). Greenspace will provide communities with accessible and safe environments to exercise, socialize, and overcome activity restriction in the COVID-19 era. Turfgrass and associated greenspace enhance the creative, intellectual, and cognitive skills of children (Frumkin, 2001; Heerwagen & Orians, 2002; Kahn & Kellert, 2002) and provide safe playing surfaces for athletes (Mack et al., 2019). According to the Centers for Disease Control and Prevention, people with close access to parks and trails are

more likely to lead a more physically active lifestyle and therefore have reduced risk of chronic diseases (Barrett, Miller, & Frumkin, 2014; CDC, 2014) and lower body mass index (BMI) (Bell, Wilson, & Liu, 2008; Liu et al., 2007). Decreased mortality rates have been documented among those living in areas with increased greenspace (James, Hart, Banay, & Laden, 2016). In addition to physical health benefits, turfgrasses also offer many mental health benefits (Moore, 1981). Proximity to greenspace and connection with nature provides a sense of tranquility (Frumkin, 2001) and has been associated with reducing stress and symptoms of depression (Barrett et al., 2014; Beyer et al., 2014), conditions likely to rise in response to COVID-19 (Gao et al., 2020). Increased participation in outdoor activities has been documented in the COVID-19 era (Venter et al., 2020), in part because exposure risks are lower outdoors (Ratnesar-Shumate et al., 2020). Turfgrasses also provide a sense of communal space for neighborhood gatherings and community events, which helps improve social ties, community pride, and overall quality of life (Hartig, Mitchell, de Vries, & Frumkin, 2014).

2.1 | Golf courses

Environmental benefits of golf courses include flood control and groundwater recharge of filtered water when absorbing surface flows from residential areas (Bouwer, 2002; Dai et al., 2016). Biodiversity and abundance of wildlife and insects are maintained as many golf courses provide the minimum size necessary for multiple species, which is critical as populations decline globally (Beninde, Veith, & Hochkirch, 2015; Tanner and Gange, 2005; van Klink et al., 2020). Golf course rough and non-play areas make up 60–70% of most golf courses and often consist of conservatively managed grasses, trees, and shrubs. When combined with the open areas of fairways and greens, this greenspace creates viable wildlife habitat (Cohn, 2008). Golf courses, like parks and forests, reduce the urban heat island effect; in Los Angeles, CA, temperatures on the golf course were $\sim 4^\circ\text{C}$ cooler than the urban surroundings (Davis, Jung, Pijanowski, & Minor, 2016; Dousset & Gourmelon, 2003). In Colorado, well-managed golf courses fairways were found to have twice as much soil carbon, up to about 80 Mg ha^{-1} , as the native prairie (Bandaranayake, Qian, Parton, Ojima, & Follett, 2003). Golf courses can be built on reclaimed landfills, providing positive environmental and social benefits (Deegan, 2017; Sharma et al. 2007).

Social benefits include the physical and mental health accrued through both playing the game of golf and the relaxation provided by being in nature, which has been shown to increase life expectancy (Barton, Hine, & Pretty,

2009; Farahmand, Broman, De Faire, Vågerö, & Ahlbom, 2009; Stenner et al., 2016). Two-thirds of U.S. states deemed golf courses as “essential services” during the COVID-19 pandemic (Kelleher, 2020). Golf courses add $\geq 25\%$ property value to homes depending on proximity (Nicholls and Crompton, 2007). The golf industry employs approximately 2 million people, with an economic impact of nearly \$180 billion (SRI, 2012). Golf courses consistently serve as fundraising venues, raising about \$4 million annually in the United States (NGF, 2017).

2.2 | Playing fields

As the majority of turfgrass playing surfaces exist in urban environments, the World Health Organization suggested that the availability, accessibility, quality, and security of public green spaces is an indicator of a healthy city (WHO, 2012). The provision, regulating, and supporting ecosystem services provided by turfgrass playing surfaces are consistent with large managed natural vegetative surfaces in urban environments (Thompson and Kao-Kniffen, 2017). Therefore, the ecosystem services provided by turfgrass playing surfaces could be explained by the preference of humans for turfgrass environments and the restorative value of organized play.

Running in a park is associated with a more restorative experience when compared to the same exercise in an urban environment (Bodin & Hartig, 2003). Bagot, Louise Allen, and Toukhsati (2015) reported that turfgrass playing fields were preferred over synthetic surfaces. Findlay and Copeland (2008) found that organized play was positively related to indices of positive adjustment and has a unique protective role for shy children by reducing anxiety. Flouri, Midouhas, and Joshi (2014) found that economically disadvantaged children in urban areas with more turfgrass playing fields had fewer emotional problems from 3 to 5 years old than their counterparts in areas with fewer turfgrass fields. To maximize ecosystem services, turfgrass playing fields should be managed to meet safety standards outlined in ASTM F-2269 and ASTM F-2060 (ASTM, 2018a, 2018b). Safety standards ensure proper care and management of turfgrass playing surfaces. Pest management is of particular concern on sports fields as failure to control infestations reduces safety to end users (Bartholomew, Campbell, & Wallace, 2015; Brosnan, Dickson, Sorochan, Thoms, & Stier, 2014).

2.3 | Parks and recreation

Parks and recreational areas are dominated by turfgrass (Ignatieva, Eriksson, Eriksson, Berg, & Hedblom, 2017;

Wheeler et al., 2017). These areas provide a variety of social, economic, and environmental benefits. The social benefits of these public areas include improved health and safer environments (Barnes et al., 2018; Demuzere et al., 2014; Dymont & Bell, 2008; Monteiro, 2017; Sadler et al., 2017). Managed turfgrass in parks serve some of the largest user groups, employ a significant number of people, and have a substantial economic impact (Diemer, 2004; English, Menard, Jensen, Brosnan, & Boyer, 2015; Hodges, Haydu, van Blokland, & Bell, 1994; Kane & Wolfe, 2012). Municipal turfgrass areas typically receive less-intensive cultural inputs and are managed with little or no pesticides (Barnes et al., 2018; Dernoeden, Carroll, & Krouse, 1994; Diesburg et al., 1997; Kowalewski et al., 2016; Kowalewski, Schwartz, Grimshaw, McCrimmon, & Layton, 2014; Patton et al., 2017; Watkins et al., 2014). Municipal turfgrass areas also serve as disposal sites for municipal waste. For example, effluent wastewater is used as an effective alternative to potable water to maintain turfgrass (Alshammary & Qian, 2008; Hayes, Mancino, & Pepper, 1990; Hyde, 1937; Mancino and Pepper, 1992; Miyamoto and Chacon, 2004; Riper and Geselbracht, 2020; Wang et al., 2014). Several cities actively promote mulching tree leaves and redistributing leaf mulch as a soil amendment that improves turfgrass health (City of Irving, 2020; City of Madison, 2020; City of Raleigh, 2020; Kowalewski et al., 2010). These waste products offer low-cost irrigation and fertilizer alternatives that are readily available to public municipalities (Heckman & Kluchinski, 1996; Riper and Geselbracht, 2020).

2.4 | Lawn and landscape

Urban vegetation and greenspaces play a crucial role in creating a comfortable living environment, particularly in semi-arid and arid climates. A study conducted in Phoenix, AZ, documented nearly 25 °C surface cooling on turfgrass compared with bare soil (Jenerette et al., 2011). Cooling from urban lawns via evapotranspiration plays a significant role in lowering surrounding surface temperatures, thereby positively affecting human thermal comfort and reducing energy consumption associated with air conditioning (Wang et al., 2016b). Amani-Beni et al. (2018) suggested increasing turfgrass coverage grown in conjunction with trees and focusing on irrigation management in order to maximize the cooling effect of urban parks.

Numerous authors have established a positive relationship between greenspaces and the well-being of residents. The restorative effect of greenspaces on humans and associated impacts on human health have been documented (Akpınar, 2016; Wells and Rollings, 2012). Akpınar, Barbosa-Leiker, and Brooks (2016) reported that more urban greenspace was associated with fewer days of

mental health complaints in urban areas. Studies also demonstrated that greenspaces in schoolyards had a higher positive restorative effect on children (Bagot et al., 2015; Kelz, Evans, & Röderer, 2015). Ward et al. (2016) and Benjamin-Neelon et al. (2019) confirmed that greenspace exposure was positively associated with physical activity in children aged 3–14 years. Significant associations between childhood obesity and the distance between the child's residence and greenspace have been reported (Manandhar et al., 2019; Shradda et al., 2019), with expansion of green areas thought to prevent weight gain at early ages.

Elam and Stigarll (2012) estimated a 17% increase in residential house pricing when landscape quality is improved. Conway, Li, Wolch, Kahle, and Jerrett (2010) documented a significant impact on house prices if homes are near a greenspace. They calculated that a 1% increase in greenspace within 200–300 ft of the residence would result in an approximate increase of 0.07% in the expected sales price. They further suggested that greening of urban areas could provide a valuable policy instrument for elevating depressed housing markets.

Studies have shown that lawns act as a net carbon sink (Braun & Bremer, 2019; Hamido et al., 2016; Law and Patton, 2017) and sequester carbon into the organic pool at similar levels to other grassland soils (Braun & Bremer, 2019). Wang et al. (2016a) documented that organic carbon sharply increased in desert soils after desert shrubs were converted to irrigated lawn. Similarly, soil inorganic carbon doubled in 6 yr in the turfgrass soil after conversion from desert shrubs.

3 | CONCLUSION


The ecosystem services provided by greenspaces containing turfgrass (i.e., golf courses, playing fields, parks and recreation, and lawn and landscapes) are predicated on the sites receiving appropriate resources for management. In the absence of allocable resources, these greenspaces will deteriorate, leading to concomitant reductions in aesthetics, function, and recreational quality. Turfgrass management encompasses a range of activities, including cultural practices (e.g., mowing, fertilization, irrigation, cultivation, and pest management), for establishing and sustaining sites at a desired level of quality.

Although budget reductions are a likely reality of the COVID-19 era, prioritization of expenditures is necessary, and essential minimums should focus on the three primary cultural practices: mowing, fertilization, and irrigation (Soldat et al., 2020). Mowing is the most basic practice needed to provide desirable turfgrass. Second, turfgrasses, like all living organisms, require nutrition. Nutrition for greenspaces primarily comes in the form of fertilizer and

other practices such as cultivation and liming to maintain soil health. Third, all plants, including turfgrasses, require water to sustain life. Irrigation resources should supplement natural rainfall, allowing these greenspaces to remain productive.

In an era of COVID-19-related budget constraints, decision makers should work with facility managers and stakeholders to identify essential minimums to ensure that realistic expectations for greenspaces are achieved while mitigating negative impacts on ecosystem services. For example, nonprioritized reductions in spending on golf course management will result in poor playing conditions, further resulting in potential revenue loss as golfers seek other venues. The cascading effect can lead to the demise of a golf course and resultant loss of valuable greenspace nestled into urban and suburban development. Similarly, insufficient resource allocation to playing fields, parks, and recreational areas can yield unsafe conditions, exposing municipalities to increased liability. Unkept and neglected landscapes are associated with higher rates of crime (Troy et al., 2016). Neglect of these greenspaces, or eliminating their inputs altogether, is not advisable.

ORCID

James T. Brosnan  <https://orcid.org/0000-0002-3390-1740>

Alec Kowalewski  <https://orcid.org/0000-0002-2854-3483>

Bernd Leinauer  <https://orcid.org/0000-0002-3700-2005>

John C. Stier  <https://orcid.org/0000-0003-0861-1095>

J. Bryan Unruh  <https://orcid.org/0000-0003-3621-7861>

REFERENCES

- Akpınar, A. (2016). How is high school greenness related to students' restoration and health? *Urban Forestry & Urban Greening*, 16, 1–8.
- Akpınar, A., Barbosa-Leiker, C., & Brooks, K. R. (2016). Does green space matter? Exploring relationships between green space type and health indicators. *Urban Forestry & Urban Greening*, 20, 407–418.
- Alshammary, S. F., & Qian, Y. L. (2008). Long term effects of effluent water irrigation on soil nitrate and phosphorus profiles under turfgrass. *Journal of Applied Sciences*, 8, 3662–3668.
- Amani-Beni, M., Zhang, B., Gao-di, X., & Jie, X. (2018). Impact of urban park's tree, grass, and waterbody on microclimate in hot summer days: A case study of Olympic Park in Beijing, China. *Urban Forestry & Urban Greening*, 32, 1–6.
- ASTM. (2018a). *Standard guide for maintaining cool season turfgrasses on athletic fields* (ASTM F2060-00). West Conshohocken, PA: ASTM International.
- ASTM. (2018b). *Standard guide for maintaining warm season turfgrasses on athletic fields* (ASTM F2269-11). West Conshohocken, PA: ASTM International.
- Bagot, K. L., Louise Allen, F. C., & Toukhsati, S. (2015). Perceived restorativeness of children's school playground environments: Nature, playground features, and play period experiences. *Journal of Environmental Psychology*, 41, 1–9.
- Baker, S. R., Bloom, N., Davis, S. J., & Terry, S. J. (2020). *COVID-induced economic uncertainty* (Working paper no. 26983). Cambridge, MA: National Bureau of Economics Research. <https://doi.org/10.3386/w26983>
- Bandaranayake, W., Qian, Y. L., Parton, W. J., Ojima, D. S., & Follett, R. F. (2003). Estimation of soil organic carbon changes in turfgrass systems using the CENTURY model. *Agronomy Journal*, 95, 558–563.
- Barnes, M. R., Nelson, K. C., Meyer, A. J., Watkins, E., Bonos, S. A., Horganc, B. P., ... Yue, C. (2018). Public land managers and sustainable urban vegetation: The case of low-input turfgrasses. *Urban Forestry and Urban Greening*, 29, 284–292.
- Barrett, M. A., Miller, D., & Frumkin, H. (2014). Parks and health: Aligning incentives to create innovations in chronic disease prevention. *Preventing Chronic Disease*, 11, 130407. <https://doi.org/10.5888/pcd11.130407>
- Bartholomew, C., Campbell, B. L., & Wallace, V. (2015). Factors affecting school grounds and athletic field quality after pesticide bans: The case of Connecticut. *HortScience*, 50, 99–103.
- Barton, J., Hine, R., & Pretty, J. (2009). The health benefits of walking in greenspaces of high natural and heritage value. *Journal of Integrative Environmental Sciences*, 6, 261–278.
- Beard, J. B., & Green, R. L. (1994). The role of turfgrasses in environmental protection and their benefits to humans. *Journal of Environmental Quality*, 23, 452–460. <https://doi.org/10.2134/jeq1994.00472425002300030007x>
- Beard, J. B., & Johns, D. (1985). The comparative heat dissipation from three typical urban surfaces: Asphalt, concrete, and bermudagrass turf. In *Texas turfgrass research 1985* (PR-4329, pp. 125–133). College Station: Texas Agricultural Experiment Station.
- Bell, J. F., Wilson, J. S., & Liu, G. C. (2008). Neighborhood greenness and 2-year changes in body mass index of children and youth. *American Journal of Preventive Medicine*, 35, 547–533.
- Beninde, J., Veith, M., & Hochkirch, A. (2015). Biodiversity in cities needs space: A meta-analysis of factors determining intra-urban biodiversity variation. *Ecology Letters*, 18, 581–592.
- Benjamin-Neelon, S. E., Platt, A., Bacardi-Gascon, M., Armstrong, S., Neelon, B., & Jimenez-Cruz, A. (2019). Greenspace, physical activity, and BMI in children from two cities in northern Mexico. *Preventive Medicine Reports*, 14, 100870. <https://doi.org/10.1016/j.pmedr.2019.100870>
- Beyer, K. M. M., Kaltenbach, A., Szabo, A., Bogar, S., Nieto, F. J., & Malecki, K. M. (2014). Exposure to neighborhood green space and mental health: Evidence from the survey of the health of Wisconsin. *International Journal of Environmental Research and Public Health*, 11, 3453–3472.
- Bodin, M., & Hartig, T. (2003). Does the outdoor environment matter for psychological restoration gained through running? *Psychology of Sport and Exercise*, 4, 141–153.
- Bouwer, H. (2002). Integrated water management for the 21st century: Problems and solutions. *Journal of Irrigation and Drainage Engineering*, 128, 193–202.
- Braun, R. C., & Bremer, D. J. (2019). Carbon sequestration in zoysiagrass turf under different irrigation and fertilization management regimes. *Agrosystems, Geosciences & Environment*, 2, 180060. <https://doi.org/10.2134/age2018.12.0060>
- Brosnan, J. T., Dickson, K. H., Sorochan, J. C., Thoms, A. W., & Stier, J. C. (2014). Large crabgrass, white clover, and hybrid bermudagrass

- athletic field playing quality in response to simulated traffic. *Crop Science*, 54, 1838–1843.
- Centers for Disease Control and Prevention (CDC). (2014). *Healthy places: Parks, trails, and health*. Retrieved from <https://www.cdc.gov/healthyplaces/healthtopics/parks.htm>
- Christians, N. E., Patton, A. J., & Law, Q. D. (2017). *Fundamentals of turfgrass management*. Hoboken, NJ: John Wiley and Sons.
- City of Irving. (2020). Leaf management. Irving, TX. Retrieved from <https://www.cityofirving.org/614/Leaf-Management>
- City of Madison. (2020). Make leaf mulch. Retrieved from <https://www.cityofmadison.com/streets/LeaveTheLeaf/mulch.cfm>
- City of Raleigh. (2020). Yard waste center. Retrieved from <https://raleighnc.gov/services/content/SolidWaste/Articles/YardWasteRecyclingCen.html>
- Cohn, J. P. (2008). Wildlife scores on golf courses. *Bioscience*, 58, 568. <https://doi.org/10.1641/B580617>
- Conway, D., Li, C. Q., Wolch, J., Kahle, C., & Jerrett, M. (2010). A spatial autocorrelation approach for examining the effects of urban greenspace on residential property values. *Journal of Real Estate Finance and Economics*, 41(2), 150–169. <https://doi.org/10.1007/s11146-008-9159-6>
- Crouch, J. A., Carter, Z., Ismaiel, A., & Roberts, J. A. (2017). The US National Mall microbiome: A census of rhizosphere bacteria inhabiting landscape turf. *Crop Science*, 57(Suppl. 1), S-341–S-348. <https://doi.org/10.2135/cropsci2016.10.0849>
- Dai, Z., Puyang, X., & Han, L. (2016). Using assessment of net ecosystem services to promote sustainability of golf course in China. *Ecological Indicators*, 63, 165–171.
- Davis, A. Y., Jung, J., Pijanowski, B. C., & Minor, E. S. (2016). Combined vegetation volume and “greenness” affect urban air temperature. *Applied Geography*, 71, 106–114.
- Deegan, J. S. (2017). *Grass over garbage: Golf courses give landfill sites a second life*. Retrieved from <https://www.golfadvisor.com/articles/grass-over-garbage-golf-courses-give-landfill-sites-a-second-life>
- Demuzere, M., Orru, K., Heidrich, O., Olazabal, E., Geneletti, D., Orru, H., ... Faehnle, M. (2014). Mitigating and adapting to climate change: Multi-functional and multi-scale assessment of green urban infrastructure. *Journal of Environmental Management*, 146, 107–115.
- Dernoeden, P. H., Carroll, M. J., & Krouse, J. M. (1994). Mowing of three fescue species for low-maintenance turf sites. *Crop Science*, 34, 1645–1649.
- Diesburg, K. L., Christians, N. E., Moore, R., Branham, B., Danneberger, T. K., Reicher, Z., ... Newman, R. (1997). Species for low-input sustainable turf in the U.S. Upper Midwest. *Agronomy Journal*, 89, 690–696.
- Diemer, J. (2004). *Economic impact: Golf and turfgrass in New Mexico 2004–2005*. Las Cruces: Department of Economics and Agricultural Business New Mexico State University.
- Dousset, B., & Gourmelon, F. (2003). Satellite multi-sensor data analysis of urban surface temperatures and landcover. *Journal of Photogrammetry and Remote Sensing*, 58, 43–54.
- Dyment, J. E., & Bell, A. C. (2008). Grounds for movement: Green school grounds as sites for promoting physical activity. *Health Education Research*, 23, 952–962.
- Emmett, E., & Stigarll, A. (2012). Landscape and house appearance impacts on the price of single-family houses. *Journal of Environmental Horticulture*, 30, 182–188. <https://doi.org/10.24266/0738-2898.30.4.182>
- English, B. C., Menard, J., Jensen, K., Brosnan, J. T., & Boyer, C. (2015). *Tennessee's turfgrass industry: Economic contributions, problematic issues, and perceptions of future growth, 2015*. Knoxville: Agri-Industry Modeling and Analysis Group University of Tennessee.
- Farahmand, B., Broman, G., De Faire, U., Vågerö, D., & Ahlbom, A. (2009). Golf: A game of life and death—Reduced mortality in Swedish golf players. *Scandinavian Journal of Medicine & Science in Sports*, 19, 419–424.
- Findlay, L., & Coplan, R. J. (2008). Come out and play: Shyness in childhood and the benefits of organized sports participation. *Canadian Journal of Behavioural Science*, 40(3), 153–161.
- Flouri, E., Midouhas, E., & Joshi, H. (2014). The role of urban neighbourhood green space in children's emotional and behavioural resilience. *Journal of Environmental Psychology*, 40, 179–186. <https://doi.org/10.1016/j.jenvp.2014.06.007>
- Frumkin, H. (2001). Beyond toxicity: Human health and the natural environment. *American Journal of Preventive Medicine*, 20, 234–240. [https://doi.org/10.1016/S0749-3797\(00\)00317-2](https://doi.org/10.1016/S0749-3797(00)00317-2)
- Gao, J., Zheng, P., Jia, Y., Chen, H., Mao, Y., Chen, S., ... Dai, J. (2020). Mental health problems and social media exposure during COVID-19 outbreak. *PLOS ONE*, 15(4), e0231924. <https://doi.org/10.1371/journal.pone.0231924>
- Grant, C. D., Campbell, C. J., & Charnock, N. R. (2002). Selection of species suitable for derelict mine site rehabilitation in New South Wales, Australia. *Water Air Soil Pollution*, 139, 215–235. <https://doi.org/10.1023/A:1015860025136>
- Hamido, S. A., Guertal, E. A., & Wood, C. W. (2016). Carbon sequestration under warm season turfgrasses in home lawns. *Journal of Geoscience and Environment Protection*, 9, 53–63. <https://doi.org/10.4236/gep.2016.49005>
- Hartig, T., Mitchell, R., de Vries, S., & Frumkin, H. (2014). Nature and health. *Annual Review of Public Health*, 35, 21.1–21.22.
- Hayes, A. R., Mancino, C. F., & Pepper, I. L. (1990). Irrigation of turfgrass with secondary sewage effluent: I. Soil and leachate water quality. *Agronomy Journal*, 82, 939–943.
- Heckman, J. R., & Kluchinski, D. (1996). Chemical composition of municipal leaf waste and hand-collected urban leaf litter. *Journal of Environmental Quality*, 25, 355–362. <https://doi.org/10.2134/jeq1996.00472425002500020021x>
- Heerwagen, J. H., & Orians, G. H. (2002). The ecological world of children. In P. H. J. Kahn & S. R. Kellert (Eds.), *Children and nature: Psychological, sociocultural, and evolutionary investigations* (pp. 29–64). Cambridge MA: MIT Press.
- Hodges, A. W., Haydu, J. J., van Blokland, P. J., & Bell, A. P. (1994). *Contributions of the turfgrass industry to Florida's economy, 1991–92: A value-added approach*. Gainesville: Institute of Food and Agricultural Sciences, University of Florida.
- Hyde, C. G. (1937). The beautification and irrigation of golden gate park with activated sludge effluent. *Sewage Works Journal*, 9, 929–941.
- Ignatieva, M., Eriksson, F., Eriksson, T., Berg, P., & Hedblom, M. (2017). The lawn as a social and cultural phenomenon in Sweden. *Urban Forestry and Urban Greening*, 21, 213–223.
- James, P., Hart, J. E., Banay, R. F., & Laden, F. (2016). Exposure to greenness and mortality in a nationwide prospective cohort study of women. *Environmental Health Perspectives*, 124, 1344–1352.
- Jenerette, G. D., Harlan, S. L., Stefanov, W. L., & Martin, C. A. (2011). Ecosystem services and urban heat riskscape moderation: Water,

- green spaces, and social inequality in Phoenix, USA. *Ecological Applications*, 21(7), 2637–2651.
- Kahn, Jr., P. H., & Kellert, S. R. (2002). *Children and nature: Psychological, sociocultural, and evolutionary investigations*. Cambridge MA: MIT Press.
- Kane, S. P., & Wolfe, K. L. (2012). *Economic contribution of turfgrass production, ornamental horticulture, landscape services, and related industry in the Georgia economy, 2010* (pp. 1–16). Athens: Center for Agribusiness and Economic Development, The University of Georgia.
- Kelleher, S. R. (2020, April 10). Are golf courses essential during the COVID-19 crisis? It depends where you live. *Forbes*. Retrieved from <https://www.forbes.com/sites/suzannerowankelleher/2020/04/10/are-golf-courses-essential-during-covid-19-crisis-it-depends-where-you-live/#54f5522a1a24>
- Kelz, C., Evans, G. W., & Röderer, K. (2015). The restorative effects of redesigning the schoolyard: A multi-methodological, quasi-experimental study in rural Austrian middle schools. *Environment & Behavior*, 47(2), 119–139
- Kowalewski, A. R., Calhoun, R. N., Hathaway, A. D., & Rogers, J. N. (2010). Using cultural practices and leaf mulch to control weeds in established turfgrass. *Applied Turfgrass Science*, 7, 1–10.
- Kowalewski, A. R., Schwartz, B. M., Grimshaw, A. L., McCrimmon, J. N., & Layton, J. M. (2014). Mowing requirement and cost to maintain bermudagrass is influenced by cultivar selection and trinexapac-ethyl use. *Applied Turfgrass Science*, 11, 1–9.
- Kowalewski, A. R., Stock, T. W., McDonald, B. W., Mattox, C. M., & Davis court, B. L. (2016). Developing and delivering a needs-based integrated pest management program for public school grounds employees. *Natural Science Education*, 45, 1–14.
- Krebs, C. C. (2020). *Advisory memorandum on identification of essential critical infrastructure workers during COVID-19 response*. U.S. Department of Homeland Security. Retrieved from https://www.cisa.gov/sites/default/files/publications/Version_3.0_CISA_Guidance_on_Essential_Critical_Infrastructure_Workers_3.pdf
- Krishnan, G., Horst, G. L., & Shea, P. J. (2000). Differential tolerance of cool- and warm-season grasses to TNT-contaminated soil. *International Journal of Phytoremediation*, 2, 369–382. <https://doi.org/10.1080/15226510008500045>
- Kuo, F. E., & Sullivan, W. C. (2001). Environment and crime in the inner city: Does vegetation reduce crime? *Environment and Behavior*, 33, 343–367.
- Kuo, F. E., Sullivan, W. C., Coley, R. L., & Brunson, L. (1998). Fertile ground for community: Inner-city neighborhood common spaces. *American Journal of Community Psychology*, 26(6), 823–851
- Laverne, R. J., & Winson-Geideman, K. (2003). The influence of trees and landscaping on rental rates at office buildings. *Arboricultural Journal*, 29(5), 281–290.
- Law, Q. D., & Patton, A. J. (2017). Biogeochemical cycling of carbon and nitrogen in cool-season turfgrass systems. *Urban Forestry & Urban Greening*, 26, 158–162.
- Lin, C. H., Lerch, R. N., Garre, H. E., & George, M. F. (2004). Incorporating forage grasses in riparian buffers for bioremediation of atrazine, isoxaflutole, and nitrate in Missouri. *Agroforestry Systems*, 63, 91–99. <https://doi.org/10.1023/B:AGFO.0000049437.70313.ef>
- Liu, G. C., Wilson, J. S., Qi, R., & Ying, J. (2007). Green neighborhoods, food retail, and childhood overweight: Differences by population density. *American Journal of Health Promotion*, 21(4S), 317–325.
- Lusk, J., Anderson, J., Charlton, D., Coble, K., Davis, A., Dewey, A., ... Zilberman, D. (2020, June). *CAST Commentary: Economic impacts of COVID-19 on food and agricultural markets*. Retrieved from <https://www.cast-science.org/wp-content/uploads/2020/06/QTA2020-3-COVID-Impacts.pdf>
- Mack, C. D., Hershman, E. B., Anderson, R. B., Coughlin, M. J., McNitt, A. S., Sendor, R. R., & Kent, R. W. (2019). Higher rates of lower extremity injury on synthetic turf compared with natural turf among National Football League athletes: Epidemiologic confirmation of a biomechanical hypothesis. *American Journal of Sports Medicine*, 47(1), 189–196. <https://doi.org/10.1177/0363546518808499>.
- Manandhar, S., Suksaroj, T. T., & Rattanapan, C. (2019). The association between green space and the prevalence of overweight/obesity among primary school children. *International Journal of Occupational and Environmental Medicine*, 10, 1–10. <http://doi.org/10.15171/ijoem.2019.1425>
- Mancino, C. F., & Pepper, I. L. (1992). Irrigation of turfgrass with secondary sewage effluent: Soil quality. *Agronomy Journal*, 84, 650–654.
- Miyamoto, S., & Chacon, A. (2004). Soil salinity of urban turf areas irrigated with saline water: II. Soil factors. *Landscape and Urban Planning*, 77, 28–38.
- Monteiro, J. A. (2017). Ecosystem services from turfgrass landscapes. *Urban Forestry and Greening*, 26, 151–157. <https://doi.org/10.1016/j.ufug.2017.04.001>
- Montgomery, J. A., Klimas, C. A., Arcus, J., DeKnock, C., Rico, K., Rodriguez, Y., ... Williams, A. (2016). Soil quality assessment is a necessary first step for designing urban green infrastructure. *Journal of Environmental Quality*, 45(1), 18–25.
- Moore, P. (1981). The well-managed lawn. *Nature*, 291, 14. <https://doi.org/10.1038/291014a0>
- National Golf Foundation (NGF). (2017). *Estimating the charitable impact of golf*. Jupiter, FL: National Golf Foundation.
- National Golf Foundation (NGF). (2020). *COVID-19 update*. Retrieved from <https://www.thengfq.com/covid-19/>
- Nicholls, S., & Crompton, J. L. (2007). The impact of a golf course on residential property values. *Journal of Sport Management*, 21, 555–570.
- Parmet, W. E., & Sinha, M. S. (2020). Covid-19: The law and limits of quarantine. *New England Journal of Medicine*, 382, e28. <https://doi.org/10.1056/NEJMp2004211>
- Patton, A. J., Schwartz, B. M., & Kenworthy, K. E. (2017). Zoysiagrass (*Zoysia* spp.) history, utilization, and improvement in the United States: A review. *Crop Science*, 57, 37–72.
- Qian, Y. L., & Follett, R. F. (2002). Assessing soil carbon sequestration in turfgrass systems using long-term soil testing data. *Agronomy Journal*, 94, 930–935.
- Ratnesar-Shumate, S., Williams, G., Green, B., Krause, M., Holland, B., Wood, S., ... Dabisch, P. (2020). Simulated sunlight rapidly inactivates SARS-CoV-2 on surfaces. *Journal of Infectious Diseases*, 222(2), 214–222. <http://doi.org/10.1093/infdis/jiaa274>
- Riper, C. V., & Geselbracht, J. (2020). Water reclamation and reuse. *Water Environment Research*, 7, 720–728.
- Sadler, R. C., Pizarro, J., Turchan, B., Gasteyer, S. P., & McGarrell, E. F. (2017). Exploring the spatial-temporal relationships between a

- community greening program and neighborhood rates of crime. *Applied Geography*, 83, 13–26.
- Shahandeh, H., & Hossner, L. R. (2000). Plant screening for chromium phytoremediation. *International Journal of Phytoremediation*, 2, 31–51. <https://doi.org/10.1080/15226510008500029>
- Sharma, H. D., & De, A. (2007). Municipal solid waste landfill settlement: Postclosure perspectives. *Journal of Geotechnical and Geoenvironmental Engineering*, 133, 619–629.
- Shraddha, M., Suksaroj, T. T., & Rattanapan, C. (2019). The association between green space and the prevalence of overweight/obesity among primary school children. *International Journal of Occupational and Environmental Medicine*, 10(1), 1–10. <http://doi.org/10.15171/ijoem.2019.1425>
- Sohrabi, C., Alsafi, Z., O'Neil, N., Khan, M., Kerwan, A., Al-Jabir, A., ... Agha, R. (2020). World Health Organization declares a global emergency: A review of the 2019 novel coronavirus (COVID-19). *International Journal of Surgery*, 76, 71–76.
- Soldat, D. J., Brosnan, J. T., Chandra, A., Gaussoin, R. E., Kowalewski, A., Leinauer, B., ... Unruh, J. B. (2020). Estimating economic minimums of mowing, fertilizing, and irrigating turfgrass. *Agricultural & Environmental Letters*, e220032. <https://doi.org/10.1002/ael2.20032>
- SRI International. (2012). *The 2011 golf economy report*. Retrieved from <http://wearegolf.org/wp-content/uploads/2019/01/2011-golf-econ-exec-sum-sri-final-12-17-12.pdf>
- State of Michigan. (2020). Coronavirus Executive Order 2020–42 FAQs. Retrieved from https://www.michigan.gov/coronavirus/0,9753,7-406-98178_98455-525278-,00.html
- State of Oregon. (2020). Masks required statewide. Retrieved from <https://govstatus.egov.com/or-oha-face-coverings>
- Stenner, B. J., Mosewich, A. D., & Buckley, J. D. (2016). An exploratory investigation into the reasons why older people play golf. *Qualitative Research in Sport, Exercise and Health*, 8, 257–272.
- Stier, J. C., Steinke, K., Ervin, E. H., Higginson, F. R., & McMaugh, P. E. (2013). Turfgrass benefits and issues. In J. C. Stier, B. P. Horgan, & S. A. Bonos (Eds.), *Turfgrass: Biology, use, and management*. Madison, WI: ASA.
- Tanner, R. A., & Gange, A. C. (2005). Effects of golf courses on local biodiversity. *Landscape Urban Planning*, 71, 137–146.
- Thwaites, R. H., Ashmore, M. R., Morton, A. J., & Pakeman, R. J. (2006). The effects of tropospheric ozone on the species dynamics of calcareous grassland. *Environmental Pollution*, 144, 500–509.
- Thompson, G. L., & Kao-Kniffin, J. (2017). Applying biodiversity and ecosystem function theory to turfgrass management. *Crop Science*, 57, S-238–S-248. <https://doi.org/10.2135/cropsci2016.05.0433>
- Thompson, O. A., Wolf, D. C., Mattice, J. D., & Thoma, G. J. (2008). Influence of nitrogen addition and plant root parameters on phytoremediation of pyrene-contaminated soil. *Water, Air, and Soil Pollution*, 189, 37–47. <https://doi.org/10.1007/s11270-007-9552-4>
- Tordoff, G. M., Baker, A. J. M., & Willis, A. J. (2000). Current approaches to the revegetation and reclamation of metalliferous mine wastes. *Chemosphere*, 41, 219–228. [https://doi.org/10.1016/S00456535\(99\)00414-2](https://doi.org/10.1016/S00456535(99)00414-2)
- Troy, A., Nunery, A., & Grove, J. M. (2016). The relationship between residential yard management and neighborhood crime: An analysis from Baltimore City and County. *Landscape and Urban Planning*, 147, 78–87.
- van Klink, R., Bowler, D. E., Gongalsky, K. B., Swengel, A. B., Gentile, A., & Chase, J. M. (2020). Meta-analysis reveals declines in terrestrial but increases in freshwater insect abundances. *Science*, 386, 417–420.
- Van Renterghem, T. V., Forssen, J., Attenborough, K., Jean, P., Defrance, J., Hornikx, M., & Kang, J. (2015). Using natural means to reduce surface transport noise during propagation outdoors. *Applied Acoustics*, 92, 86–101. <https://doi.org/10.1016/j.apacoust.2015.01.004>
- Venter, Z., Barton, D., Gundersen, V., Figari, H., & Nowell, M. (2020). Urban nature in a time of crisis: Recreational use of green space increases during the COVID-19 outbreak in Oslo, Norway. *Environmental Research Letters*, <https://doi.org/10.1088/1748-9326/abb396>
- Wang, F., Qiao, M., Lv, Z., Guo, G., Jia, Y., Su, Y., & Zhu, Y. (2014). Impact of reclaimed water irrigation on antibiotic resistance in public parks, Beijing, China. *Environmental Pollution*, 184, 247–253.
- Wang, J., Monger, C., Wang, X., Serena, M., & Leinauer, B. (2016a). Carbon sequestration in response to grassland-shrubland-turfgrass conversions and a test for carbonate biomineralization in desert soils, New Mexico, USA. *Soil Science Society of America Journal*, 80, 1591–1603.
- Wang, Z., Zhao, X., Yang, J., & Song, J. (2016b). Cooling and energy saving potentials of shade trees and urban lawns in a desert city. *Applied Energy*, 161, 437–444.
- Ward, J. S., Duncan, J. S., Jarden, A., & Stewart, T. (2016). The impact of children's exposure to greenspace on physical activity, cognitive development, emotional wellbeing, and ability to appraise risk. *Health & Place*, 40, 44–50. <https://doi.org/10.1016/j.healthplace.2016.04.015>
- Watkins, E., Gardner, D. S., Stier, J. C., Soldat, D. J., St. John, R. A., Christians, N. E., ... Gaussoin, R. E. (2014). Cultivar performance of low-input turfgrass species for the north central United States. *Applied Turfgrass Science*, 11, 1–7.
- Wheeler, M. M., Neill, C., Groffman, P. M., Avolio, M., Bettez, N., Cavender-Bares, J., ... Trammell, T. L. E. (2017). Continental-scale homogenization of residential lawn plant communities. *Landscape and Urban Planning*, 165, 54–63.
- Wells, N. M., & Rollings, K. (2012). The natural environment in residential settings: Influences on human health and function. In *The Oxford handbook of environmental and conservation psychology* (pp. 509–523). Oxford, U.K.: Oxford University Press.
- World Health Organization (WHO). (2020). Coronavirus disease (COVID-19) dashboard. Retrieved from https://covid19.who.int/?gclid=Cj0KCQjwnv71BRCOARIsAikxW9HQzUwWD8F3oyzv0L_92a9hH9FqaamwdG5DnPP5SX2hKV5Xbf7pkekaAn1bEALw_wcB
- World Health Organization (WHO). (2012). Health indicators of sustainable cities. In *Proceedings of the United Nations Conference on Sustainable Development, Rio de Janeiro, Brazil, 20–22 June 2012*.

How to cite this article: Brosnan JT, Chandra A, Gaussoin RE, et al. A justification for continued management of turfgrass during economic contraction. *Agric Environ Lett*. 2020;5:1–8. <https://doi.org/10.1002/ael2.20033>